Chapter - 7 Integrals

edukalpclasses.com

Miscellaneous Exercise

Question 1: $\frac{1}{x-x^3}$ Answer 1: $\frac{1}{x-x^3} = \frac{1}{x(1-x^2)} = \frac{1}{x(1-x)(1+x)}$ Let $\frac{1}{x(1-x)(1+x)} = \frac{A}{x} + \frac{B}{(1-x)} + \frac{C}{1+x}$...(1) $\Rightarrow 1 = A(1-x^2) + Bx(1+x) + Cx(1-x)$ $\Rightarrow 1 = A - Ax^2 + Bx + Bx^2 + Cx - Cx^2 + C$ Equating the coefficients of x^2 , x, and constant term, we obtain -A + B - C = 0B + C = 0A = 1On solving these equations, we obtain $A = 1, B = \frac{1}{2}$, and $C = -\frac{1}{2}$ From equation (1), we obtain $\frac{1}{x(1-x)(1+x)} = \frac{1}{x} + \frac{1}{2(1-x)} - \frac{1}{2(1+x)}$ $\Rightarrow \int \frac{1}{x(1-x)(1+x)} dx = \int \frac{1}{x} dx + \frac{1}{2} \int \frac{1}{1-x} dx - \frac{1}{2} \int \frac{1}{1+x} dx$ $= \log |x| - \frac{1}{2} \log |(1-x)| - \frac{1}{2} \log |(1+x)|$ $= \log |x| - \log \left| (1-x)^{\frac{1}{2}} \right| - \log \left| (1+x)^{\frac{1}{2}} \right|$ $= \log \left| \frac{x}{(1-x)^{\frac{1}{2}}(1+x)^{\frac{1}{2}}} \right| + C$ $= \log \left| \left(\frac{x^2}{1 - x^2} \right)^{\frac{1}{2}} \right| + C$ $=\frac{1}{2}\log\left|\frac{x^2}{1-x^2}\right|+C$

For more study Materials login to edukalpclasses.com

1

Chapter - 7 Integrals

Question 2:

$$\frac{1}{\sqrt{x+a} + \sqrt{(x+b)}}$$

Answer 2:



Question 3:

1		а
$\frac{1}{\sqrt{a_{1}}}$		x = -
$x \sqrt{ax - x}$	[Hint: Put	1

Answer 3:



Question 4:

$$\frac{1}{x^2\left(x^4+1\right)^{\frac{3}{4}}}$$

Answer 4:

$$\frac{1}{x^2\left(x^4+1\right)^{\frac{3}{4}}}$$

Multiplying and dividing by x^{-3} , we obtain

$$\frac{x^{-3}}{x^2 \cdot x^{-3} (x^4 + 1)^{\frac{3}{4}}} = \frac{x^{-3} (x^4 + 1)^{\frac{3}{4}}}{x^2 \cdot x^{-3}}$$

$$= \frac{(x^4 + 1)^{\frac{3}{4}}}{x^5 \cdot (x^4)^{\frac{3}{4}}}$$

$$= \frac{1}{x^5} (\frac{x^4 + 1}{x^4})^{\frac{3}{4}}$$

$$= \frac{1}{x^5} (\frac{x^4 + 1}{x^4})^{\frac{3}{4}}$$

$$= \frac{1}{x^5} (1 + \frac{1}{x^4})^{\frac{3}{4}}$$

$$= \frac{1}{x^5} (x^4 + 1)^{\frac{3}{4}} dx = dt \Rightarrow \frac{1}{x^5} dx = -\frac{dt}{4}$$

$$\therefore \int \frac{1}{x^2 (x^4 + 1)^{\frac{3}{4}}} dx = \int \frac{1}{x^5} (1 + \frac{1}{x^4})^{-\frac{3}{4}} dx$$

$$= -\frac{1}{4} \int (1 + t)^{\frac{3}{4}} dt$$

$$= -\frac{1}{4} \left[\frac{(1 + t)^{\frac{1}{4}}}{\frac{1}{4}} \right] + C$$

$$= -\frac{1}{4} \left[\frac{(1 + t)^{\frac{1}{4}}}{\frac{1}{4}} \right] + C$$

$$= -\left[(1 + \frac{1}{x^4})^{\frac{1}{4}} + C \right]$$



Answer 5:



Chapter - 7 Integrals

edukalpclasses.com

Question 6:

$$\frac{5x}{(x+1)(x^2+9)}$$

Answer 6:

Let
$$\frac{5x}{(x+1)(x^2+9)} = \frac{A}{(x+1)} + \frac{Bx+C}{(x^2+9)}$$
 ...(1)
 $\Rightarrow 5x = A(x^2+9) + (Bx+C)(x+1)$
 $\Rightarrow 5x = Ax^2 + 9A + Bx^2 + Bx + Cx + C$

ational .

Equating the coefficients of x^2 , x, and constant term, we obtain

$$A + B = 0 B$$

+ C = 5
 $9A + C = 0$

On solving these equations, we obtain

$$A = -\frac{1}{2}, B = \frac{1}{2}, \text{ and } C = \frac{9}{2}$$

From equation (1), we obtain

$$\frac{5x}{(x+1)(x^2+9)} = \frac{-1}{2(x+1)} + \frac{x+\frac{9}{2}}{(x^2+9)}$$
EET

$$\int \frac{5x}{(x+1)(x^2+9)} dx = \int \left\{ \frac{-1}{2(x+1)} + \frac{(x+9)}{2(x^2+9)} \right\} dx$$

$$= -\frac{1}{2} \log|x+1| + \frac{1}{2} \int \frac{x}{x^2+9} dx + \frac{9}{2} \int \frac{1}{x^2+9} dx$$

$$= -\frac{1}{2} \log|x+1| + \frac{1}{4} \int \frac{2x}{x^2+9} dx + \frac{9}{2} \int \frac{1}{x^2+9} dx$$

$$= -\frac{1}{2} \log|x+1| + \frac{1}{4} \log|x^2+9| + \frac{9}{2} \cdot \frac{1}{3} \tan^{-1} \frac{x}{3}$$

$$= -\frac{1}{2} \log|x+1| + \frac{1}{4} \log(x^2+9) + \frac{3}{2} \tan^{-1} \frac{x}{3} + C$$

Question 7: $\sin x$ $\sin(x-a)$ Answer 7: $\sin x$ $\sin(x-a)$ Let $x - a = t \Rightarrow dx = dt$ $\int \frac{\sin x}{\sin (x-a)} dx = \int \frac{\sin (t+a)}{\sin t} dt$ $= \int \frac{\sin t \cos a + \cos t \sin a}{\sin t} dt$ $= \int (\cos a + \cot t \sin a) dt$ $= t \cos a + \sin a \log |\sin t| + C_1$ $= (x-a)\cos a + \sin a \log |\sin (x-a)| + C_1$ $= x \cos a + \sin a \log \left| \sin \left(x - a \right) \right| - a \cos a + C_1$ $= \sin a \log |\sin (x-a)| + x \cos a + C$ **Question 8:** $e^{5\log x} - e^{4\log x}$ $e^{3\log x} - e^{2\log x}$ Answer 8: $\frac{e^{5\log x} - e^{4\log x}}{e^{3\log x} - e^{2\log x}} = \frac{e^{4\log x} \left(e^{\log x} - 1\right)}{e^{2\log x} \left(e^{\log x} - 1\right)}$ $=e^{2\log x}$ $=e^{\log x^2}$ $=x^2$ $\therefore \int \frac{e^{5\log x} - e^{4\log x}}{e^{3\log x} - e^{2\log x}} dx = \int x^2 dx = \frac{x^3}{3} + C$

Question 9:

 $\frac{\cos x}{\sqrt{4-\sin^2 x}}$

 $\sqrt{4-\sin x}$

Answer 9:

$$\frac{\cos x}{\sqrt{4-\sin^2 x}}$$

Let sin x = t \Rightarrow cos x dx = dt



Question 11:

$$\frac{1}{\cos(x+a)\cos(x+b)}$$

Answer 11:

$$\frac{1}{\cos(x+a)\cos(x+b)}$$
Multiplying and dividing by $\sin(a-b)$, we obtain
$$\frac{1}{\sin(a-b)} \left[\frac{\sin(a-b)}{\cos(x+a)\cos(x+b)} \right]$$

$$= \frac{1}{\sin(a-b)} \left[\frac{\sin[(x+a)-(x+b)]}{\cos(x+a)\cos(x+b)} \right]$$

$$= \frac{1}{\sin(a-b)} \left[\frac{\sin(x+a)-\cos(x+b)-\cos(x+a)\sin(x+b)}{\cos(x+a)\cos(x+b)} \right]$$

$$= \frac{1}{\sin(a-b)} \left[\frac{\sin(x+a)}{\cos(x+a)} - \frac{\sin(x+b)}{\cos(x+a)} \right]$$

$$= \frac{1}{\sin(a-b)} \left[\tan(x+a) - \tan(x+b) \right]$$

$$\int \frac{1}{\cos(x+a)\cos(x+b)} dx = \frac{1}{\sin(a-b)} \int \left[\tan(x+a) - \tan(x+b) \right] dx$$

$$= \frac{1}{\sin(a-b)} \left[-\log|\cos(x+a)| + \log|\cos(x+b)| \right] + C$$



Question 14:

 $\frac{1}{\left(x^2+1\right)\left(x^2+4\right)}$

Answer 14:

$$\therefore \frac{1}{(x^2+1)(x^2+4)} = \frac{Ax+B}{(x^2+1)} + \frac{Cx+D}{(x^2+4)}$$
$$\Rightarrow 1 = (Ax+B)(x^2+4) + (Cx+D)(x^2+1)$$
$$\Rightarrow 1 = Ax^3 + 4Ax + Bx^2 + 4B + Cx^3 + Cx + Dx^2 + Dx^2$$

tional

Equating the coefficients of x^3 , x^2 , x, and constant term, we obtain

A + C = 0 B + D = 0 4A + C = 04B + D = 1

On solving these equations, we obtain

$$A = 0, B = \frac{1}{3}, C = 0, \text{and } D = -\frac{1}{3}$$

From equation (1), we obtain

$$\frac{1}{(x^2+1)(x^2+4)} = \frac{1}{3(x^2+1)} - \frac{1}{3(x^2+4)}$$
$$\int \frac{1}{(x^2+1)(x^2+4)} dx = \frac{1}{3} \int \frac{1}{x^2+1} dx - \frac{1}{3} \int \frac{1}{x^2+4} dx$$
$$= \frac{1}{3} \tan^{-1} x - \frac{1}{3} \cdot \frac{1}{2} \tan^{-1} \frac{x}{2} + C$$
$$= \frac{1}{3} \tan^{-1} x - \frac{1}{6} \tan^{-1} \frac{x}{2} + C$$

Chapter - 7 Integrals

edukalpclasses.com





Question 19:

$$\frac{\sin^{-1}\sqrt{x} - \cos^{-1}\sqrt{x}}{\sin^{-1}\sqrt{x} + \cos^{-1}\sqrt{x}}, x \in [0,1]$$

Answer 19:

Let
$$I = \int \frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} dx$$

It is known that, $\sin^{-1}\sqrt{x} + \cos^{-1}\sqrt{x} = \frac{\pi}{2}$ $\Rightarrow I = \int \frac{\left(\frac{\pi}{2} - \cos^{-1}\sqrt{x}\right) - \cos^{-1}\sqrt{x}}{\frac{\pi}{2}} dx \text{ cational Servic}$ $=\frac{2\pi}{\pi}\left(\frac{1}{2}-2\cos^{-1}\sqrt{x}\right)dx$ $=\frac{2\pi}{\pi}\frac{1}{2}\int 1 \cdot dx - \frac{4}{\pi}\int \cos^{-1}\sqrt{x} \, dx$ $=x-\frac{4}{\pi}\int\cos^{-1}\sqrt{x}\,dx$ Let $I_1 = \int \cos^{-1} \sqrt{x} \, dx$ Also, let $\sqrt{x} = t \implies dx = 2t dt$ $\Rightarrow I_1 = 2 \int \cos^{-1} t \cdot t \, dt$ $= 2 \left| \cos^{-1} t \cdot \frac{t^2}{2} - \int \frac{-1}{\sqrt{1-t^2}} \cdot \frac{t^2}{2} dt \right|$ $=t^2\cos^{-1}t+\int \frac{t^2}{\sqrt{1-t^2}}dt$ $=t^{2}\cos^{-1}t - \int \frac{1-t^{2}-1}{\sqrt{1-t^{2}}}dt$ $=t^{2}\cos^{-1}t - \int \sqrt{1-t^{2}}dt + \int \frac{1}{\sqrt{1-t^{2}}}dt$ $= t^{2} \cos^{-1} t - \frac{t}{2} \sqrt{1 - t^{2}} - \frac{1}{2} \sin^{-1} t + \sin^{-1} t$ $=t^{2}\cos^{-1}t - \frac{t}{2}\sqrt{1-t^{2}} + \frac{1}{2}\sin^{-1}t$

From equation (1), we obtain



edukalpclasses.com



$$= -4 \int \sin^2 \frac{\theta}{2} \cos \theta \, d\theta$$

$$= -4 \int \sin^2 \frac{\theta}{2} \cdot \left(2 \cos^2 \frac{\theta}{2} - 1\right) d\theta$$

$$= -4 \int \left(2 \sin^2 \frac{\theta}{2} \cos^2 \frac{\theta}{2} - \sin^2 \frac{\theta}{2}\right) d\theta$$

$$= -8 \int \sin^2 \frac{\theta}{2} \cdot \cos^2 \frac{\theta}{2} \, d\theta + 4 \int \sin^2 \frac{\theta}{2} \, d\theta$$

$$= -2 \int \sin^2 \theta \, d\theta + 4 \int \sin^2 \frac{\theta}{2} \, d\theta$$

$$= -2 \int \left(\frac{1 - \cos 2\theta}{2}\right) d\theta + 4 \int \frac{1 - \cos \theta}{2} \, d\theta$$

$$= -2 \left[\frac{\theta}{2} - \frac{\sin 2\theta}{4}\right] + 4 \left[\frac{\theta}{2} - \frac{\sin \theta}{2}\right] + C$$

$$= -\theta + \frac{\sin 2\theta}{2} + 2\theta - 2\sin \theta + C$$

$$= \theta + \frac{2\sin \theta \cos \theta}{2} - 2\sin \theta + C$$

$$= \theta + \frac{2\sin \theta \cos \theta}{2} - 2\sin \theta + C$$

$$= \theta + \sqrt{1 - \cos^2 \theta} \cdot \cos \theta - 2\sqrt{1 - \cos^2 \theta} + C$$

$$= -2\sqrt{1 - x} + \cos^{-1} \sqrt{x} + \sqrt{x(1 - x)} + C$$

$$= -2\sqrt{1 - x} + \cos^{-1} \sqrt{x} + \sqrt{x - x^2} + C$$

Question 21: $\frac{2+\sin 2x}{1+\cos 2x}e^x$ Answer 21: $I = \int \left(\frac{2 + \sin 2x}{1 + \cos 2x}\right) e^x$ $= \int \left(\frac{2+2\sin x \cos x}{2\cos^2 x}\right) e^x$ $=\int \left(\frac{1+\sin x \cos x}{\cos^2 x}\right) e^x$ $= \int (\sec^2 x + \tan x)e^x$ Let $f(x) = \tan x \Rightarrow f'(x) = \sec^2 x$ $\therefore I = \int (f(x) + f'(x)] e^{x} dx$ $=e^{x}f(x)+C$ $=e^{x} \tan x + C$ **Question 22:** $x^2 + x + 1$ $\overline{(x+1)^2(x+2)}$ Answer 22: Let $\frac{x^2 + x + 1}{(x+1)^2(x+2)} = \frac{A}{(x+1)} + \frac{B}{(x+1)^2} + \frac{C}{(x+2)}$ 1....(1) $\Rightarrow x^{2} + x + 1 = A(x+1)(x+2) + B(x+2) + C(x^{2}+2x+1)$ $\Rightarrow x^{2} + x + 1 = A(x^{2} + 3x + 2) + B(x + 2) + C(x^{2} + 2x + 1)$ $\Rightarrow x^2 + x + 1 = (A+C)x^2 + (3A+B+2C)x + (2A+2B+C)$ Equating the coefficients of x^2 , x and constant term, we obtain A + C = 1

3A + B + 2C = 1 2A + 2B + C = 1On solving these equations, we obtain A = -2, B = 1, and C = 3From equation (1), we obtain

edukalpclasses.com

$$\frac{x^{2} + x + 1}{(x+1)^{2}(x+2)} = \frac{-2}{(x+1)} + \frac{3}{(x+2)} + \frac{1}{(x+1)^{2}}$$
$$\int \frac{x^{2} + x + 1}{(x+1)^{2}(x+2)} dx = -2 \int \frac{1}{x+1} dx + 3 \int \frac{1}{(x+2)} dx + \int \frac{1}{(x+1)^{2}} dx$$
$$= -2 \log|x+1| + 3 \log|x+2| - \frac{1}{(x+1)} + C$$

Question 23: $\tan^{-1}\sqrt{\frac{1-x}{1+x}}$ Answer 23: $I = \tan^{-1} \sqrt{\frac{1-x}{1+x}} dx$ Let $x = \cos\theta \implies dx = -\sin\theta d\theta$ $I = \int \tan^{-1} \sqrt{\frac{1 - \cos\theta}{1 + \cos\theta}} \left(-\sin\theta d\theta \right)$ $= -\int \tan^{-1} \sqrt{\frac{2\sin^2\frac{\theta}{2}}{2\cos^2\frac{\theta}{2}}} \sin\theta d\theta}$ $=-\int \tan^{-1} \tan \frac{\theta}{2} \cdot \sin \theta d\theta$ $=-\frac{1}{2}\int \theta \cdot \sin \theta d\theta$ $= -\frac{1}{2} \Big[\theta \cdot (-\cos\theta) - \int 1 \cdot (-\cos\theta) d\theta \Big]$ $=-\frac{1}{2}\left[-\theta\cos\theta+\sin\theta\right]$ $=+\frac{1}{2}\theta\cos\theta-\frac{1}{2}\sin\theta$ $=\frac{1}{2}\cos^{-1}x \cdot x - \frac{1}{2}\sqrt{1-x^2} + C$ $=\frac{x}{2}\cos^{-1}x - \frac{1}{2}\sqrt{1-x^2} + C$ $=\frac{1}{2}\left(x\cos^{-1}x-\sqrt{1-x^{2}}\right)+C$

For more study Materials login to edukalpclasses.com

18

Chapter - 7 Integrals

Question 24:

$$\frac{\sqrt{x^2+1} \left[\log \left(x^2+1\right) - 2 \log x \right]}{x^4}$$

Answer 24:



Integrating by parts, we obtain

edukalpclasses.com



For more study Materials login to edukalpclasses.com

20

Chapter - 7 Integrals

edukalpclasses.com



For more study Materials login to edukalpclasses.com

21

Question 27:

 $\int_0^{\frac{\pi}{2}} \frac{\cos^2 x \, dx}{\cos^2 x + 4\sin^2 x}$

Answer 27:

Let
$$I = \int_{0}^{\frac{\pi}{2}} \frac{\cos^{2} x}{\cos^{2} x + 4\sin^{2} x} dx$$

$$\Rightarrow I = \int_{0}^{\frac{\pi}{2}} \frac{\cos^{2} x}{\cos^{2} x + 4(1 - \cos^{2} x)} dx$$

$$\Rightarrow I = \int_{0}^{\frac{\pi}{2}} \frac{\cos^{2} x}{\cos^{2} x + 4 - 4\cos^{2} x} dx$$

$$\Rightarrow I = \int_{0}^{\frac{\pi}{2}} \int_{0}^{\frac{\pi}{2}} \frac{4 - 3\cos^{2} x}{4 - 3\cos^{2} x} dx$$

$$\Rightarrow I = \frac{-1}{3} \int_{0}^{\frac{\pi}{2}} \frac{4 - 3\cos^{2} x}{4 - 3\cos^{2} x} dx + \frac{1}{3} \int_{0}^{\frac{\pi}{2}} \frac{4 - 3\cos^{2} x}{4 - 3\cos^{2} x} dx$$

$$\Rightarrow I = \frac{-1}{3} \int_{0}^{\frac{\pi}{2}} 1 dx + \frac{1}{3} \int_{0}^{\frac{\pi}{2}} \frac{4 \sec^{2} x}{4 \sec^{2} x - 3} dx$$

$$\Rightarrow I = \frac{-1}{3} \int_{0}^{\frac{\pi}{2}} 1 dx + \frac{1}{3} \int_{0}^{\frac{\pi}{2}} \frac{4 \sec^{2} x}{4 \sec^{2} x - 3} dx$$

$$\Rightarrow I = \frac{-1}{3} [x]_{0}^{\frac{\pi}{2}} + \frac{1}{3} \int_{0}^{\frac{\pi}{2}} \frac{4 \sec^{2} x}{4 \sec^{2} x - 3} dx$$

$$\Rightarrow I = -\frac{\pi}{6} + \frac{2}{3} \int_{0}^{\frac{\pi}{2}} \frac{2 \sec^{2} x}{1 + 4 \tan^{2} x} dx$$
Let $2 \tan x = t \Rightarrow 2 \sec^{2} x dx$

$$= \int_{0}^{\infty} \frac{dt}{1 + t^{2}}$$

$$\Rightarrow \int_{0}^{\frac{\pi}{2}} \frac{2 \sec^{2} x}{1 + 4 \tan^{2} x} dx = \int_{0}^{\infty} \frac{dt}{1 + t^{2}}$$

$$= [\tan^{-1} t]_{0}^{\infty}$$

$$= [\tan^{-1} t]_{0}^{\infty}$$

Therefore, from (1), we obtain

$$I = -\frac{\pi}{6} + \frac{2}{3} \left[\frac{\pi}{2} \right] = \frac{\pi}{3} - \frac{\pi}{6} = \frac{\pi}{6}$$

Question 28:

$$\int_{a}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$$
Answer 28:
Let $I = \int_{\frac{\pi}{6}}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$
 $\Rightarrow I = \int_{a}^{\frac{\pi}{3}} \frac{(\sin x + \cos x)}{\sqrt{-(-\sin 2x)}} dx$
 $\Rightarrow I = \int_{a}^{\frac{\pi}{3}} \frac{\sin x + \cos x}{\sqrt{-(-1+1-2\sin x \cos x)}} dx$
 $\Rightarrow I = \int_{a}^{\frac{\pi}{3}} \frac{(\sin x + \cos x)}{\sqrt{1-(\sin^2 x + \cos^2 x - 2\sin x \cos x)}} dx$
 $\Rightarrow I = \int_{a}^{\frac{\pi}{3}} \frac{(\sin x + \cos x) dx}{\sqrt{1-(\sin x - \cos x)^2}}$
Let $(\sin x - \cos x) = t \Rightarrow (\sin x + \cos x) dx = dt$
When $x = \frac{\pi}{6}, t = (\frac{1-\sqrt{3}}{2})$ and when $x = \frac{\pi}{3}, t = (\frac{\sqrt{3}-1}{2})$
 $I = \int_{-\frac{\sqrt{3}-1}{2}}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}}$
 $\Rightarrow I = \int_{-\frac{\sqrt{3}-1}{2}}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^2}}$
As $\frac{1}{\sqrt{1-(-t)^2}} = \frac{1}{\sqrt{1-t^2}}$, therefore, $\frac{1}{\sqrt{1-t^2}}$ is an even function.
 $\int_{a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx$

It is known that if f(x) is an even function, then

$$\Rightarrow I = 2 \int_{0}^{\frac{\sqrt{3}-1}{2}} \frac{dt}{\sqrt{1-t^{2}}} \\ = \left[2\sin^{-1} t \right]_{0}^{\frac{\sqrt{3}-1}{2}} \\ = 2\sin^{-1} \left(\frac{\sqrt{3}-1}{2} \right)$$

Chapter - 7 Integrals

Question 29:

$$\int_0^1 \frac{dx}{\sqrt{1+x} - \sqrt{x}}$$

Answer 29:

Let
$$I = \int_{0}^{1} \frac{dx}{\sqrt{1+x} - \sqrt{x}}$$

 $I = \int_{0}^{1} \frac{1}{(\sqrt{1+x} - \sqrt{x})} \times \frac{(\sqrt{1+x} + \sqrt{x})}{(\sqrt{1+x} + \sqrt{x})} dx$
 $= \int_{0}^{1} \frac{\sqrt{1+x} + \sqrt{x}}{1+x-x} dx$
 $= \int_{0}^{1} \sqrt{1+x} dx + \int_{0}^{1} \sqrt{x} dx$
 $= \left[\frac{2}{3}(1+x)^{\frac{3}{2}}\right]_{0}^{1} + \left[\frac{2}{3}(x)^{\frac{3}{2}}\right]_{0}^{1}$
 $= \frac{2}{3}\left[(2)^{\frac{3}{2}} - 1\right] + \frac{2}{3}[1]$
 $= \frac{2}{3}(2)^{\frac{3}{2}}$
 $= \frac{4\sqrt{2}}{3}$

Question 30:

$$\int_0^{\frac{\pi}{4}} \frac{\sin x + \cos x}{9 + 16\sin 2x} dx$$

Answer 30:

Let
$$I = \int_{0}^{\pi} \frac{\sin x + \cos x}{9 + 16 \sin 2x} dx$$

Also, let $\sin x - \cos x = t \Rightarrow (\cos x + \sin x) dx = dt$
When $x = 0$, $t = -1$ and when $x = \frac{\pi}{4}$, $t = 0$
 $\Rightarrow (\sin x - \cos x)^{2} = t^{2}$
 $\Rightarrow \sin^{2} x + \cos^{2} x - 2 \sin x \cos x = t^{2}$
 $\Rightarrow 1 - \sin 2x = t^{2}$
 $\Rightarrow \sin 2x = 1 - t^{2}$
 $\therefore I = \int_{-1}^{0} \frac{dt}{9 + 16(1 - t^{2})}$
 $= \int_{-1}^{0} \frac{dt}{9 + 16 - 16t^{2}}$
 $= \int_{-1}^{0} \frac{dt}{25 - 16t^{2}} = \int_{-1}^{0} \frac{dt}{(5)^{2} - (4t)^{2}}$
 $= \frac{1}{4} \left[\frac{1}{2(5)} \log \left| \frac{5 + 4t}{5 - 4t} \right| \right]_{-1}^{0}$
 $= \frac{1}{40} \left[\log(1) - \log \left| \frac{1}{9} \right| \right]$

Question 31:

 $\int_0^{\frac{\pi}{2}} \sin 2x \tan^{-1} \left(\sin x \right) dx$

Answer 31:

Let
$$I = \int_{0}^{\frac{\pi}{2}} \sin 2x \tan^{-1} (\sin x) dx = \int_{0}^{\frac{\pi}{2}} 2\sin x \cos x \tan^{-1} (\sin x) dx$$

Also, let $\sin x = t \Rightarrow \cos x dx = dt$
When $x = 0, t = 0$ and when $x = \frac{\pi}{2}, t = 1$ **tion all Solutions**
 $\Rightarrow I = 2\int_{0}^{1} t \tan^{-1} t dt = \tan^{-1} t \cdot \int t dt - \int \left\{ \frac{d}{dt} (\tan^{-1} t) \int t dt \right\} dt$
 $= \tan^{-1} t \cdot \frac{t^{2}}{2} - \int \int \frac{1}{1+t^{2}} \frac{t^{2}}{2} dt$
 $= \frac{t^{2} \tan^{-1} t}{2} - \int \frac{1}{2} \int \frac{t^{2} + 1 - 1}{1+t^{2}} dt$
 $= \frac{t^{2} \tan^{-1} t}{2} - \frac{1}{2} \int t dt + \frac{1}{2} \int \frac{1}{1+t^{2}} dt$
 $= \frac{t^{2} \tan^{-1} t}{2} - \frac{1}{2} \int t dt + \frac{1}{2} \int \frac{1}{1+t^{2}} dt$
 $= \frac{t^{2} \tan^{-1} t}{2} - \frac{1}{2} \cdot t + \frac{1}{2} \tan^{-1} t$
 $\Rightarrow \int_{0}^{1} t \cdot \tan^{-1} t dt = \left[\frac{t^{2} \cdot \tan^{-1} t}{2} - \frac{t}{2} + \frac{1}{2} \tan^{-1} t \right]_{0}^{1}$
 $= \frac{1}{2} \left[\frac{\pi}{4} - 1 + \frac{\pi}{4} \right]$
 $= \frac{1}{2} \left[\frac{\pi}{2} - 1 \right] = \frac{\pi}{4} - \frac{1}{2}$

From equation (1), we obtain

$$I = 2\left[\frac{\pi}{4} - \frac{1}{2}\right] = \frac{\pi}{2} - 1$$

Chapter - 7 Integrals

edukalpclasses.com

Question 32:

 $\int_0^{\pi} \frac{x \tan x}{\sec x + \tan x} dx$

Answer 32:

Let
$$I = \int_{0}^{\pi} \frac{x \tan x}{\sec x + \tan x} dx$$
 ...(1)
 $I = \int_{0}^{\pi} \left\{ \frac{(\pi - x) \tan (\pi - x)}{\sec (\pi - x) + \tan (\pi - x)} \right\} dx$ $\left(\int_{0}^{\pi} f(x) dx = \int_{0}^{\pi} f(a - x) dx \right\}$
 $\Rightarrow I = \int_{0}^{\pi} \left\{ \frac{-(\pi - x) \tan x}{-(\sec x + \tan x)} \right\} dx$
Adding (1) and (2), we obtain
 $2I = \int_{0}^{\pi} \frac{\pi \tan x}{\sec x + \tan x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{\cos x}{\cos x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{\cos x}{1 + \sin x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{\sin x}{1 + \sin x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{1 - \sin x}{\cos x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{1 - \sin x}{\cos^{2} x} dx$
 $\Rightarrow 2I = \pi \int_{0}^{\pi} \frac{1 - \sin x}{\cos^{2} x} dx$
 $\Rightarrow 2I = \pi \left[x \right]_{0}^{\pi} - \pi \int_{0}^{\pi} \frac{1 - \sin x}{\cos^{2} x} dx$
 $\Rightarrow 2I = \pi^{2} - \pi \left[\tan x - \sec x \right]_{0}^{\pi} \frac{1 - \sin x}{\cos^{2} x} dx$
 $\Rightarrow 2I = \pi^{2} - \pi \left[\tan x - \sec x \right]_{0}^{\pi} \frac{1 - \sin x}{\cos^{2} x} dx$
 $\Rightarrow 2I = \pi^{2} - \pi \left[\tan \pi - \sec x - \tan 0 + \sec 0 \right]$
 $\Rightarrow 2I = \pi^{2} - \pi \left[0 - (-1) - 0 + 1 \right]$
 $\Rightarrow 2I = \pi^{2} - \pi \left[0 - (-1) - 0 + 1 \right]$

Question 33:

$$\int_{0}^{4} \left[|x-1| + |x-2| + |x-3| \right] dx$$

Answer 33:

Let
$$I = \int_{1}^{4} [|x-1|| + |x-2| + |x-3|] dx$$

 $\Rightarrow I = \int_{1}^{4} |x-1| dx + \int_{1}^{4} |x-2| dx + \int_{1}^{4} |x-3| dx$
 $I = I_{1} + I_{2} + I_{3}$...(1)
where, $I_{1} = \int_{1}^{4} |x-1| dx$, $I_{2} = \int_{1}^{4} |x-2| dx$, and $I_{3} = \int_{1}^{4} |x-3| dx$
 $I_{1} = \int_{1}^{4} |x-1| dx$
 $(x-1) \ge 0$ for $1 \le x \le 4$
 $\therefore I_{1} = \int_{1}^{4} (x-1) dx$
 $\Rightarrow I_{1} = \left[8 - 4 - \frac{1}{2} + 1 \right] = \frac{9}{2}$...(2)
 $I_{2} = \int_{1}^{4} |x-2| dx$
 $x - 2 \ge 0$ for $2 \le x \le 4$ and $x - 2 \le 0$ for $1 \le x \le 2$
 $\therefore I_{2} = \int_{1}^{4} |x-2| dx$
 $\Rightarrow I_{2} = \left[2x - \frac{x^{2}}{2} \right]_{1}^{2} + \left[\frac{x^{2}}{2} - 2x \right]_{2}^{4}$
 $\Rightarrow I_{2} = \left[4 - 2 - 2 + \frac{1}{2} \right] + [8 - 8 - 2 + 4]$
 $\Rightarrow I_{2} = \frac{1}{2} + 2 = \frac{5}{2}$...(3)

Chapter - 7 Integrals

$$I_{3} = \int^{4} |x-3| dx$$

$$x-3 \ge 0 \text{ for } 3 \le x \le 4 \text{ and } x-3 \le 0 \text{ for } 1 \le x \le 3$$

$$\therefore I_{3} = \int^{3} (3-x) dx + \int^{4} (x-3) dx$$

$$\Rightarrow I_{3} = \left[3x - \frac{x^{2}}{2} \right]^{3} + \left[\frac{x^{2}}{2} - 3x \right]^{4}_{3}$$

$$\Rightarrow I_{3} = \left[9 - \frac{9}{2} - 3 + \frac{1}{2} \right] + \left[8 - 12 - \frac{9}{2} + 9 \right]$$

$$\Rightarrow I_{3} = \left[6 - 4 \right] + \left[\frac{1}{2} \right] = \frac{5}{2} \qquad \dots(4)$$

From equations (1), (2), (3), and (4), we obtain

$$I = \frac{9}{2} + \frac{5}{2} + \frac{5}{2} = \frac{19}{2}$$

Question 34:

Question 34:

$$\int_{x^{2}}^{3} \frac{dx}{(x+1)} = \frac{2}{3} + \log \frac{2}{3}$$
Answer 34:
Let $I = \int_{x^{2}}^{3} \frac{dx}{(x+1)}$
Also, let $\frac{1}{x^{2}(x+1)} = \frac{A}{x} + \frac{B}{x^{2}} + \frac{C}{x+1}$
 $\Rightarrow 1 = Ax(x+1) + B(x+1) + C(x^{2})$
 $\Rightarrow 1 = Ax^{2} + Ax + Bx + B + Cx^{2}$

Equating the coefficients of x^2 , x, and constant term, we obtain

$$A + C = 0$$

 $A + B =$
 0
 $B = 1$
On solving these equations, we obtain
 $A = -1, C = 1$, and $B = 1$

$$\therefore \frac{1}{x^2 (x+1)} = \frac{-1}{x} + \frac{1}{x^2} + \frac{1}{(x+1)}$$

$$\Rightarrow I = \int_1^3 \left\{ -\frac{1}{x} + \frac{1}{x^2} + \frac{1}{(x+1)} \right\} dx$$

$$= \left[-\log x - \frac{1}{x} + \log (x+1) \right]_1^3$$

$$= \left[\log \left(\frac{x+1}{x} \right) - \frac{1}{x} \right]_1^3$$

$$= \log \left(\frac{4}{3} \right) - \frac{1}{3} - \log \left(\frac{2}{1} \right) + 1$$

$$= \log 4 - \log 3 - \log 2 + \frac{2}{3}$$

$$= \log \left(\frac{2}{3} \right) + \frac{2}{3}$$

Hence, the given result is proved.

Question 35:

 $\int xe^x dx = 1$

Answer 35:

Let
$$I = \int_0^1 x e^x dx$$

Integrating by parts, we obtain

$$I = x \int_{0}^{1} e^{x} dx - \int_{0}^{1} \left\{ \left(\frac{d}{dx}(x) \right) \int e^{x} dx \right\} dx$$
$$= \left[x e^{x} \right]_{0}^{1} - \int_{0}^{1} e^{x} dx$$
$$= \left[x e^{x} \right]_{0}^{1} - \left[e^{x} \right]_{0}^{1}$$
$$= e - e + 1$$
$$= 1$$
Hence, the given result is proved

Hence, the given result is proved.

Question 36:

 $\int_{-1}^{1} x^{17} \cos^4 x \, dx = 0$

Answer 36:

Let $I = \int_{-1}^{1} x^{17} \cos^4 x dx$ Also, let $f(x) = x^{17} \cos^4 x$ $\Rightarrow f(-x) = (-x)^{17} \cos^4 (-x) = -x^{17} \cos^4 x = -f(x)$ Therefore, f (x) is an odd function.

It is known that if f(x) is an odd function, then $\int_{a}^{a} f(x) dx = 0$

:
$$I = \int_{-1}^{1} x^{17} \cos^4 x \, dx = 0$$

Hence, the given result is proved.

Question 37:

 $\int_{0}^{\frac{\pi}{2}} \sin^3 x \, dx = \frac{2}{3}$

Answer 37:

Let
$$I = \int_{0}^{\frac{\pi}{2}} \sin^{3} x \, dx$$

 $I = \int_{0}^{\frac{\pi}{2}} \sin^{2} x \cdot \sin x \, dx$
 $= \int_{0}^{\frac{\pi}{2}} (1 - \cos^{2} x) \sin x \, dx$
 $= \int_{0}^{\frac{\pi}{2}} \sin x \, dx - \int_{0}^{\frac{\pi}{2}} \cos^{2} x \cdot \sin x \, dx$
 $= [-\cos x]_{0}^{\frac{\pi}{2}} + \left[\frac{\cos^{3} x}{3}\right]_{0}^{\frac{\pi}{2}}$
 $= 1 + \frac{1}{3}[-1] = 1 - \frac{1}{3} = \frac{2}{3}$

Hence, the given result is proved.



edukalpclasses.com

Question 38:

 $\int_{4}^{\pi} 2 \tan^3 x dx = 1 - \log 2$

Answer 38:

Let
$$I = \int_{0}^{\frac{\pi}{4}} 2\tan^{3} x \, dx$$

 $I = 2 \int_{0}^{\frac{\pi}{4}} \tan^{2} x \tan x \, dx = 2 \int_{0}^{\frac{\pi}{4}} (\sec^{2} x - 1) \tan x \, dx$
 $= 2 \int_{0}^{\frac{\pi}{4}} \sec^{2} x \tan x \, dx - 2 \int_{0}^{\frac{\pi}{4}} \tan x \, dx$
 $\int \tan^{2} x \int_{0}^{\frac{\pi}{4}} x = \frac{\pi}{4}$

$$= 2\left\lfloor \frac{\tan x}{2} \right\rfloor_{0} + 2\left[\log \cos x\right]_{0}^{\overline{4}}$$
$$= 1 + 2\left[\log \cos \frac{\pi}{4} - \log \cos 0\right]$$
$$= 1 + 2\left[\log \frac{1}{\sqrt{2}} - \log 1\right]$$

 $= 1 - \log 2 - \log 1 = 1 - \log 2$

Hence, the given result is proved.

Question 39:

$$\int_{0}^{1} \sin^{-1} x \, dx = \frac{\pi}{2} - 1$$

Answer 39:

Let
$$I = \int_0^1 \sin^{-1} x \, dx$$

 $\Rightarrow I = \int_0^1 \sin^{-1} x \cdot 1 \cdot dx$

Integrating by parts, we obtain

$$I = \left[\sin^{-1} x \cdot x\right]_{0}^{1} - \int_{0}^{1} \frac{1}{\sqrt{1 - x^{2}}} \cdot x \, dx$$
$$= \left[x \sin^{-1} x\right]_{0}^{1} + \frac{1}{2} \int_{0}^{1} \frac{(-2x)}{\sqrt{1 - x^{2}}} \, dx$$

Let $1 - x^2 = t \Rightarrow -2x dx = dt$



Chapter - 7 Integrals

edukalpclasses.com

When
$$x = 0$$
, $t = 1$ and when $x = 1$, $t = 0$
 $I = \left[x \sin^{-1} x\right]_{0}^{1} + \frac{1}{2} \int_{0}^{0} \frac{dt}{\sqrt{t}}$
 $= \left[x \sin^{-1} x\right]_{0}^{1} + \frac{1}{2} \left[2\sqrt{t}\right]_{1}^{0}$
 $= \sin^{-1}(1) + \left[-\sqrt{1}\right]$
 $= \frac{\pi}{2} - 1$

Hence, the given result is proved.





Question 41:

 $\int \frac{dx}{e^x + e^{-x}}$ is equal to

A. $\tan^{-1}(e^x) + C$ B. $\tan^{-1}(e^{-x}) + C$

b.
$$\tan^{-1}(e^{-x}) + C$$

$$\mathsf{C}. \qquad \log \left(e^x - e^{-x} \right) + \mathsf{C}$$

$$\mathsf{D.} \quad \log(e^x + e^{-x}) + \mathsf{C}$$

Answer 41:

Let
$$I = \int \frac{dx}{e^x + e^{-x}} dx = \int \frac{e^x}{e^{2x} + 1} dx$$

Also, let $e^x = t \implies e^x dx = dt$
 $\therefore I = \int \frac{dt}{1 + t^2} dx$
 $= \tan^{-1} t + C$

Hence, the correct Answer is A.

is

 $= \tan^{-1}(e^x) + C$

Question 42:

$$\int \frac{\cos 2x}{\left(\sin x + \cos x\right)^2} dx$$

A.
$$\frac{-1}{\sin x + \cos x} + C$$

- B. $\log |\sin x + \cos x| + C$
- C. $\log |\sin x \cos x| + C$

D.
$$\frac{1}{\left(\sin x + \cos x\right)^2}$$
 equal to

Answer 42:

Let
$$I = \frac{\cos 2x}{(\cos x + \sin x)^2}$$

 $I = \int \frac{\cos^2 x - \sin^2 x}{(\cos x + \sin x)^2} dx$
 $= \int \frac{(\cos x + \sin x)(\cos x - \sin x)}{(\cos x + \sin x)^2} dx$
 $= \int \frac{\cos x - \sin x}{(\cos x + \sin x)} dx$
Let $\cos x + \sin x = t \Rightarrow (\cos x - \sin x) dx = dt$
 $\therefore I = \int \frac{dt}{t}$
 $= \log |t| + C$
 $= \log |c| + C$
 $=$

Let $I = \int_{a}^{b} x f(x) dx$...(1)

Chapter - 7 Integrals

edukalpclasses.com

$$I = \int_{a}^{b} (a+b-x) f(a+b-x) dx$$

$$\Rightarrow I = \int_{a}^{b} (a+b-x) f(x) dx$$

$$\Rightarrow I = (a+b) \int_{a}^{b} f(x) dx - I$$

$$\Rightarrow I + I = (a+b) \int_{a}^{b} f(x) dx$$

$$\Rightarrow 2I = (a+b) \int_{a}^{b} f(x) dx$$

$$\Rightarrow I = \left(\frac{a+b}{2}\right) \int_{a}^{b} f(x) dx$$

$$\left(\int_{a}^{b} f(x) dx = \int_{a}^{b} f(a+b-x) dx\right)$$

 $\left[Using(1) \right]$

Hence, the correct Answer is D.

tan⁻¹

Question 44:

The value of A. 1 B. 0 C. - 1 D. $\frac{\pi}{4}$

Answer 44:

Let
$$I = \int_{0}^{1} \tan^{-1} \left(\frac{2x-1}{1+x-x^{2}} \right) dx$$

 $\Rightarrow I = \int_{0}^{1} \tan^{-1} \left(\frac{x-(1-x)}{1+x(1-x)} \right) dx$
 $\Rightarrow I = \int_{0}^{1} \left[\tan^{-1} x - \tan^{-1} (1-x) \right] dx$...(1)
 $\Rightarrow I = \int_{0}^{1} \left[\tan^{-1} (1-x) - \tan^{-1} (1-1+x) \right] dx$
 $\Rightarrow I = \int_{0}^{1} \left[\tan^{-1} (1-x) - \tan^{-1} (x) \right] dx$
 $\Rightarrow I = \int_{0}^{1} \left[\tan^{-1} (1-x) - \tan^{-1} (x) \right] dx$...(2)
Adding (1) and (2), we obtain

2x - 1

1+x-x

dx

is



edukalpclasses.com

$$2I = \int_0^1 (\tan^{-1} x + \tan^{-1} (1 - x) - \tan^{-1} (1 - x) - \tan^{-1} x) dx$$

$$\Rightarrow 2I = 0$$

$$\Rightarrow I = 0$$

Hence, the correct Answer is B.

